
ISSN 1348-5059

気象庁海洋月報

Monthly Ocean Report

July 2004 (No. 139)



Climate and Marine Department

Japan Meteorological Agency

気象庁 気候・海洋気象部

平成16年7月

気象庁海洋月報は、気象庁 気候・海洋気象部 海洋気象情報室で作成しています。
本月報に関する問い合わせは、
〒100-8122 東京都千代田区大手町1-3-4 気象庁 気候・海洋気象部 海洋気象情報室まで。
電 話 (03) 3212-8341 (内線 5127、5166)
ＦＡＸ (03) 3211-3047
電子メール ocean@climar.kishou.go.jp

Monthly Ocean Report is prepared by the Office of Marine Prediction, Climate and Marine Department, Japan Meteorological Agency.

Correspondence relating to this report may be directed to :

Ippei Eguchi
Head, Office of Marine Prediction
Climate and Marine Department, Japan Meteorological Agency
1-3-4 Ote-machi, Chiyoda-ku, Tokyo 100-8122 Japan
Facsimile: +81-3-3211-3047
e-mail : ocean@climar.kishou.go.jp

表紙は、2004年7月の太平洋の月平均海面水温偏差図です。
陰影部は、平年より高温の海域を示しています。

The figure on the front cover shows the monthly mean SST anomalies for July 2004 in the Pacific Ocean.
Areas of positive anomalies are shaded.

目　次

* MGDSST (Merged satellite and *in-situ* data Global Daily SST)を使用している。

** 従来と同様の、現場観測による海面水温をもとにした解析値を使用している。

Table of Contents

* Based on MGDSST (Merged satellite and *in-situ* data Global Daily SST).

** Based on the traditional analysis using *in-situ* SSTs.

海況の特徴（２００４年７月）

日本近海の海況

海面水温は三陸沖を除いて全般に平年より高かった。能登半島沖から九州沿岸、房総半島沖から日本の南にかけて、及び北海道の東で平年より1℃以上高く、東海から紀伊半島にかけての沿岸では平年より1.5℃以上高かった（第４図)。

黒潮は、中旬までは四国の南で32° N以南を流れていたが、下旬には足摺岬に接近して流れていた。紀伊半島以東では、上旬は潮岬の南から北東に流れ、中旬は137.5° E付近を北上した。下旬は紀伊半島の南東を大きく迂回した後、138° E付近を北上した。その後、御前崎の南を東進して三宅島付近を通過し、房総半島沖を北東に流れていた（第９図)。

太平洋の海況

太平洋赤道域の海面水温は、日付変更線付近、170° W付近と160° Wから140°Wにかけて平年より1℃以上高かった。125° E付近と110°W以東では平年より0.5℃以上低かった。

北太平洋の海面水温は全般に平年より高く、35° N、165° E付近、アラスカ湾、及びベーリング海で+2℃を超える正偏差が見られた（第１７図)。

Highlights July 2004

Seas adjacent to Japan

SSTs were above normal in the seas adjacent to Japan except off Sanriku. SSTs were more than 1℃ above normal from off the Noto Peninsula to around Kyushu, from off the Boso Peninsula to south of Japan, and east of Hokkaido. SSTs were more than 1.5℃ above normal in the seas from off Tokai to the Kii Peninsula (Fig. 4).


In the seas south of Shikoku, the Kuroshio flowed south of 32°N in early and middle July, and flowed near Ashizuri Misaki in late July. It turned NE-ward off south of Shiono Misaki in early July. In middle July it flowed northward along 137.5°E. In late July, it flowed far off southeast of the Kii Peninsula and flowed northward along 138°E. After flowing eastward south of Omaezaki, it passed near Miyake Island and flowed NE-ward off the Boso Peninsula (Fig. 9).

Pacific Ocean

In the equatorial Pacific, SSTs were more than 1℃ above normal around the date line, around 170°W, and from 160°W to 140°W. SSTs were more than 0.5℃ below normal around 125°E and east of 110°W.

SSTs were generally above normal in the North Pacific. Positive SST anomalies exceeding +2℃ were found in the seas around 35°N,165°E, in the Gulf of Alaska and in the Bering Sea (Fig. 17).

Ten-day mean SST & Anomalies




旬平均沿岸水温(℃)
Coastal Water Temperature

地点名 (Index Number)	水温 Temp.	偏差 Anom.
石　垣(47918)	28.0	-0.9
御前崎(47655)	23.4	0.6
小名浜(47598)	17.2	-0.1
宮　古(47585)	16.7	2.0
江　差(47428)	18.6	0.7
浜　田(47755)	25.0	2.3

第1図 北西太平洋旬平均海面水温図(上)及び偏差図(下)(2004年 7月上旬)

Fig. 1 Ten-day mean SST (top) and anomalies (bottom) in the western North Pacific for the 1st ten-day period in July 2004. Anomalies are computed from the JMA climatology (1971-2000). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.

Ten-day mean SST & Anomalies



旬平均沿岸水温(℃)


Coastal Water Temperature

地点名 (Index Number)	水温 Temp.	偏差 Anom.
石　垣(47918)	28.6	-0.6
御前崎(47655)	23.9	0.1
小名浜(47598)	19.1	0.9
宮　古(47585)	17.1	1.5
江　差(47428)	20.7	1.7
浜　田(47755)	26.5	2.2

第2図 北西太平洋旬平均海面水温図(上)及び偏差図(下)(2004年 7月中旬)

Fig. 2 Ten-day mean SST (top) and anomalies (bottom) in the western North Pacific for the 2nd ten-day period in July 2004. Anomalies are computed from the JMA climatology (1971-2000). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.

Ten-day mean SST & Anomalies




旬平均沿岸水温(℃)
Coastal Water Temperature

地点名 (Index Number)	水温 Temp.	偏差 Anom.
石　垣(47918)	28.9	-0.3
御前崎(47655)	25.7	0.8
小名浜(47598)	19.2	-0.5
宮　古(47585)	19.0	1.9
江　差(47428)	23.1	2.2
浜　田(47755)	28.8	3.3

第3図 北西太平洋旬平均海面水温図(上)及び偏差図(下) (2004年 7月下旬)


Fig. 3 Ten-day mean SST (top) and anomalies (bottom) in the western North Pacific for the 3rd ten-day period in July 2004. Anomalies are computed from the JMA climatology (1971-2000). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.

Monthly Mean SST & Anomalies




第4図 北西太平洋月平均海面水温図(上)及び偏差図(下) (2004年 7月)

Fig. 4 Monthly mean SST (top) and anomalies (bottom) in the western North Pacific for July 2004. Anomalies are computed from the JMA climatology (1971-2000). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.



第5図　日本近海月平均海面高度分布図(上)と海面海流図(下)（2004年7月）
（海洋総合解析システムによる）


Fig. 5　Monthly mean sea surface height (top) and current (bottom) in seas adjacent to Japan for July 2004. Contour interval is 10cm (top). (calculated by the Ocean Comprehensive Analysis System)



第 6 図　日本近海月平均表層水温図（上）と表層海流図（下）（深さ 100 m）（2004 年 7 月）
（海洋総合解析システムによる）


Fig. 6　Monthly mean subsurface (100 m) temperature (top) and current (bottom) in seas adjacent to Japan for July 2004. Contour interval is 1 ℃ (top). (calculated by the Ocean Comprehensive Analysis System)

200m Temperature / Current Adjacent to Japan



第７図　日本近海月平均表層水温図（上）と表層海流図（下）　（深さ 200 m）（2004 年 7 月）
（海洋総合解析システムによる）


Fig. 7　Monthly mean subsurface (200 m) temperature (top) and current (bottom) in seas adjacent to Japan for July 2004. Contour interval is 1 ˚C (top). (calculated by the Ocean Comprehensive Analysis System)



第 8 図　日本近海月平均表層水温図（上）と表層海流図（下）（深さ 400 m）（2004 年 7 月）
（海洋総合解析システムによる）


Fig. 8　Monthly mean subsurface (400 m) temperature (top) and current (bottom) in seas adjacent to Japan for July 2004. Contour interval is 1 ˚C (top). (calculated by the Ocean Comprehensive Analysis System)

Location of the Kuroshio axis




第9図 旬別黒潮流軸図

Fig. 9 Location of the Kuroshio axis




第 10 図　北西太平洋月平均海面高度分布図 (上) と海面海流図 (下) (2004 年 7 月)
(海洋総合解析システムによる)

Fig. 10　Monthly mean sea surface height (top) and current (bottom) in the western North Pacific for July 2004. Contour interval is 10cm (top). (calculated by the Ocean Comprehensive Analysis System)



第 11 図　東経 137° 線に沿う月平均表層水温の鉛直断面図 （2004 年 7 月）
（海洋総合解析システムによる）

Fig. 11　Vertical cross section of monthly mean temperature along 137 °E for July 2004. Contour interval is 1 °C. (calculated by the Ocean Comprehensive Analysis System)




第 12 図　東経 165° 線に沿う月平均表層水温の鉛直断面図 （2004 年 7 月）
（海洋総合解析システムによる）


Fig. 12　Vertical cross section of monthly mean temperature along 165 °E for July 2004. Contour interval is 1 °C. (calculated by the Ocean Comprehensive Analysis System)



第13図　海域別旬平均海面水温平年偏差時系列図


Fig. 13 Time series of the ten-day mean SST anomalies (°C) in the western North Pacific for the areas shown in the right panel. Negative anomalies are shaded. Anomalies are computed from the JMA climatology (1971-2000).






第13図　海域別旬平均海面水温平年偏差時系列図（つづき）

Fig. 13 (continue) Time series of the ten-day mean SST anomalies (°C) in the western North Pacific for the areas shown in the right panel. Negative anomalies are shaded. Anomalies are computed from the JMA climatology (1971-2000).









第14図 沿岸水温時系列図

Fig. 14 Time series of hourly coastal water temperatures (°C) measured at 6 coastal stations of the JMA shown in the right panel.
Thin lines show ten-day mean climatology.








第15図 沿岸潮位時系列図（2004年 7月）


Fig. 15 Time series of hourly sea level observation (upper line) and its anomalies (lower line) at 9 stations of the JMA in the right panel.
Short term variations are removed from raw data.


NM: new moon, FQ: first quarter
FM: full moon, LQ: last quarter
T. P.: Mean sea level of Tokyo Bay (Tokyo Peil)













第15図 つづき
Fig. 15 continue




第16図　漂流型海洋気象ブイロボットの軌跡と観測要素の時系列図

Fig.16　Time series of JMA drifting buoy observation.
The upper panel shows paths of the buoys. Circles denote the position at 00h UTC everyday.
The lower panel shows time series of observation data.




第16図 つづき
Fig.16 continue

SST & Anomalies in the Global Ocean




第17図　全球月平均海面水温図（上）及び偏差図（下）（2004年 7月）

Fig. 17　Monthly mean SST over the global ocean (top) and anomalies (bottom) for July 2004.　Anomalies are computed from the JMA climatology (1971-2000). Contour interval is 1°C (top) and additional anomaly contours of ±0.5°C are shown by thin lines.　Negative anomalies are spotted.　Maximum coverage of sea ice is shaded.　Data sources for sea ice are U.S. National Ice Center (polar regions) and JMA (the Sea of Okhotsk).




第 18 図　太平洋月平均表層水温図（深さ 100 m）（2004 年 7 月）
（海洋総合解析システムによる）

Fig. 18　Monthly mean subsurface (100 m) temperature in the Pacific Ocean for July 2004. Contour interval is 1 ℃. (calculated by the Ocean Comprehensive Analysis System)




第 19 図　太平洋月平均表層水温図（深さ 200 m）（2004 年 7 月）
（海洋総合解析システムによる）

Fig. 19　Monthly mean subsurface (200 m) temperature in the Pacific Ocean for July 2004. Contour interval is 1 ℃. (calculated by the Ocean Comprehensive Analysis System)




第 20 図　太平洋月平均表層水温図（深さ 400 m）　(2004 年 7 月)
（海洋総合解析システムによる）

Fig. 20　Monthly mean subsurface (400 m) temperature in the Pacific Ocean for July 2004. Contour interval is 1 ℃. (calculated by the Ocean Comprehensive Analysis System)




第 21 図　太平洋ブイ流跡図（2004 年 7 月）

Fig. 21　Trajectories of drifting buoys in the Pacific Ocean during July 2004. Each track is for one buoy. Small circles represent the last positions for the month.




第22図　太平洋赤道に沿う表層水温（上）及び偏差（下）の深度－経度断面図（2004年 7月）

Fig. 22 Depth-longitude sections of temperature (top) and anomalies (bottom) along the equator in the Pacific Ocean for July 2004. Data are derived from the Ocean Data Assimilation System of JMA. Contour interval is 1°C (top) and 0.5°C (bottom). Values greater than 28°C in the top panel are shaded. In the bottom panel, values greater than 1°C are shaded and values less than -1°C are spotted. Anomalies are computed from 1987-2003 base period means.



第 23 図　太平洋月平均海面水位偏差図（2004 年 6 月）


Fig. 23　Monthly sea level anomalies for June 2004 obtained from mean sea level (1993-2001) adjusted for atmospheric pressure. This figure is prepared by the SOC of ISLP-PAC, Department of Oceanography, University of Hawaii.



第 24 図　全球月平均海面高度偏差図（2004 年 6 月）


Fig. 24　Monthly sea surface height anomalies in the global ocean derived from the altimeter data of Jason-1 for June 2004. Contour interval is 5 cm. Positive anomalies exceeding +5 cm are shaded and negative anomalies exceeding -5 cm are spotted. Anomalies are computed from 1993-1998 base period means.

Long-term Variation of Monthly Mean SST Anomalies




第25図 海域平均海面水温偏差の長期変動

Fig. 25 Long-term variations of monthly mean sea surface temperature anomalies (°C) for the areas shown in the right panel. Anomalies are computed from the JMA climatology (1971-2000). Thin lines show the monthly mean and thick lines their 13-month running means.



Time Series of Tropical Pacific SST Anomalies




第26図　太平洋熱帯域の海面水温偏差時系列図

Fig. 26 Time series of monthly mean SST anomalies (°C) in the tropical Pacific Ocean for the areas indicated at the bottom panel. Thin lines show the monthly means and thick lines their 5-month running means. Positive anomalies (A, B, C) and negative anomalies (D) are shaded. Anomalies are computed from the JMA climatology (1971-2000).

MONTH YEAR	SEA SURFACE TEMPERATURE AND ANOMALY (°C) REGION D 14N-EQ 130E-150E		REGION A 4N-4S 160E-150W		REGION B 4N-4S 150W-90W		REGION C EQ-10S 90W-80W		SOI	SEA LEVEL PRESSURE ANOMALY (hPa) TAHITI 17-33S 149-37W	DARWIN 12-24S 130-52E
JAN 02	28.7	0.2	28.7	0.7	25.0	-0.4	24.2	-0.3	0.3	1.3	0.7
FEB 02	28.6	0.3	28.5	0.5	26.3	0.0	26.7	0.7	0.7	1.5	-0.2
MAR 02	28.8	0.3	28.5	0.4	27.2	0.1	27.5	1.2	-0.2	0.4	1.0
APR 02	29.3	0.5	29.0	0.8	27.5	0.3	25.9	0.5	-0.1	0.1	0.3
MAY 02	29.7	0.4	29.4	0.8	27.2	0.3	24.9	0.6	-1.3	-1.1	0.7
JUN 02	29.8	0.4	29.5	1.0	27.0	0.8	22.4	-0.5	-0.4	-0.2	0.4
JUL 02	29.3	0.0	29.3	0.8	25.9	0.5	21.7	-0.2	-0.7	0.5	1.6
AUG 02	29.2	0.0	29.3	0.9	25.3	0.6	21.0	0.1	-1.2	-1.4	0.5
SEP 02	29.3	0.0	29.3	0.9	25.5	0.8	20.9	0.4	-0.5	-0.3	0.8
OCT 02	29.5	0.1	29.3	0.9	25.7	1.0	21.6	0.7	-0.5	-1.0	-0.3
NOV 02	29.6	0.3	29.6	1.2	26.1	1.4	22.8	1.0	-0.4	0.1	1.0
DEC 02	29.2	0.2	29.3	1.1	26.3	1.3	23.9	1.0	-1.0	-0.1	1.7
JAN 03	28.7	0.2	29.1	1.1	26.2	0.8	24.3	-0.2	-0.1	-0.1	0.2
FEB 03	28.4	0.1	28.9	0.9	26.8	0.5	26.3	0.3	-0.5	-1.5	-0.2
MAR 03	28.7	0.2	29.0	0.9	27.0	-0.1	26.5	0.2	-0.3	-0.2	0.5
APR 03	29.3	0.5	28.8	0.6	27.1	-0.1	24.9	-0.5	-0.1	0.5	0.8
MAY 03	29.7	0.4	28.9	0.3	26.0	-0.9	22.8	-1.5	-0.6	-0.7	0.2
JUN 03	29.8	0.4	29.1	0.6	25.9	-0.3	21.9	-1.0	-0.9	-0.9	0.4
JUL 03	29.6	0.3	29.0	0.5	25.6	0.2	20.9	-1.0	0.2	0.7	0.3
AUG 03	29.4	0.2	28.9	0.5	24.9	0.2	21.2	0.3	0.0	-0.2	-0.1
SEP 03	29.5	0.2	28.9	0.5	24.8	0.1	20.2	-0.3	-0.1	-0.3	0.0
OCT 03	29.7	0.3	29.1	0.7	25.2	0.5	21.1	0.2	-0.1	-0.5	-0.4
NOV 03	29.6	0.3	29.1	0.7	25.2	0.5	22.2	0.4	-0.2	0.0	0.4
DEC 03	29.2	0.2	28.9	0.7	25.5	0.5	23.4	0.5	1.1	0.9	-1.1
JAN 04	29.0	0.5	28.7	0.7	26.0	0.6	24.7	0.2	-1.0	-1.5	0.9
FEB 04	28.8	0.5	28.4	0.4	26.5	0.2	26.5	0.5	0.8	1.4	-0.6
MAR 04	29.1	0.6	28.3	0.2	27.2	0.1	26.0	-0.3	0.1	-0.7	-1.3
APR 04	29.4	0.6	28.7	0.5	27.3	0.1	25.1	-0.3	-1.4	-1.0	1.3
MAY 04	29.6	0.3	29.1	0.5	26.6	-0.3	23.3	-1.0	1.2	0.9	-0.7
JUN 04	29.4	0.0	29.1	0.6	26.2	0.0	21.9	-1.0	-1.1	0.1	1.7
JUL 04	29.6	0.3	29.3	0.8	25.4	0.0	20.6	-1.3	-0.6	-0.2	0.9

第1表　太平洋熱帯域の大気及び海洋の監視指数


Table 1 Atmospheric and oceanic monitoring indices in the tropical Pacific Ocean



第27図　赤道に沿った海面水温（左）および偏差（右）の時間-経度断面図


Fig. 27　Time-longitude sections of SSTs (left) and anomalies (right) along the equator (2°N-2°S) in the global ocean. Contour interval is 1°C (left) and 0.5°C (right). Values greater than 28°C (+0.5°C) are shaded and values less than 23°C (-0.5°C) are spotted in the left (right) panel. Dashed contours indicate negative anomalies in the right panel. Anomalies are computed from 1971-2000 base period means.

Time-Longitude Sections of Ocean Heat Content & Anomalies (Eq.)



第28図　太平洋の赤道に沿う海洋表層の貯熱量（左）及び偏差（右）の時間－経度断面図

Fig. 28 Time-longitude sections of ocean heat content in the upper 260 m along the equator (left) and anomalies (right) in the Pacific Ocean. Data are derived from the Ocean Data Assimilation System of JMA. Contour interval is 1°C (left) and 0.5°C (right). Values greater than 20°C (+0.5°C) are shaded and values less than 16°C (−0.5°C) are spotted in the left (right) panel. Dashed contours indicate negative anomalies in the right panel. Anomalies are computed from 1987-2003 base period means.




第29図　太平洋赤道域の表層水温の時間－深度断面図
（上：西部、中：中部、下：東部）

Fig. 29 Time-depth sections of equatorial subsurface temperature for the western Pacific (top, 160°E), the central Pacific (middle, 140°W) and the eastern Pacific (bottom, 110°W). Data are derived from the Ocean Data Assimilation System of JMA. Contour interval is 1°C. Values greater than 28°C are shaded.




第30-1図　北極域の海氷図（2004年 7月16日）

Fig.30-1 Sea ice chart in the Arctic on 16 Jul. 2004.
The shaded area represents the sea ice extent. The dotted line denotes the mean ice edge averaged for the period from 1972 through 1990.
The data source is the National Ice Center (USA).




第30-2図　南極域の海氷図（2004年 7月16日）

Fig.30-2 Sea ice chart in the Antarctic on 16 Jul. 2004.
The shaded area represents the sea ice extent. The dotted line denotes the mean ice edge averaged for the period from 1973 through 1990.
The data source is the National Ice Center (USA).



第30-3図 オホーツク海の海氷図（2004年 7月15日）


Fig.30-3 Sea ice chart in the Sea of Okhotsk on 15 Jul. 2004.




第30-4図 オホーツク海の海氷域面積

Fig.30-4 Time series of 5-day sea ice area in the Sea of Okhotsk.


: current season (2003/2004)
: mean extent (1971-2000)
: maximum or minimum extent (1971-2003)
: last season (2002/2003)






第 31 図 神戸海洋気象台啓風丸による東経 137 度に沿った水温鉛直断面図 (℃). (a) 水温 (2004 年 1 月), (b) 水温平年偏差 (2004 年 1 月, 平年値の統計期間は 1971～2000 年), (c) 水温 (2004 年 4 月), (d) 観測位置図.


Fig. 31 Vertical sections of water temperature (℃) along 137°E observed by R/V Keifu Maru. (a) Water temperature, (b) Water temperature anomalies in January 2004, (c) Water temperature in April 2004, and (d) Location of oceanographic observations along 137°E. Anomalies are computed from 1971–2000 base period means.






第32-1図　測点分布図(2004年 7月)

Fig. 32-1 Geographical distribution of SHIP messages (top) and BATHY/TESAC messages (bottom) for July 2004.





第32-2図　測点分布図(2004年 7月)

Fig. 32-2 Geographical distribution of BUOY messages (top) and TRACKOB messages (bottom) for July 2004.

気象庁海洋月報

Monthly Ocean Report

July 2004 (No. 139)

本月報掲載の2004年7月のプロダクトには、気象庁が観測したデータ及びGTS（全球気象通信網）を通じて入手した国際気象通報式による通報のほか、下記の機関等から提供いただいたデータを使用しています。御礼申し上げます。

防衛庁、水産庁、海上保安庁

水産総合研究センター

海洋研究開発機構

東北大学、東京大学

水産試験場等（北海道、岩手県、宮城県、山形県、福島県、茨城県、千葉県、東京都、富山県、石川県、福井県、愛知県、三重県、兵庫県、和歌山県、鳥取県、島根県、山口県、佐賀県、宮崎県）

（順不同）

平成１６年８月２０日発行

編集兼発行者　気　象　庁
東京都千代田区大手町１－３－４

印刷所　有限会社　進英プリント
東京都渋谷区代々木 1-38-19